ECOM-5170 January 1968 AD

# PREDICTING DIFFUSION OF ATMOSPHERIC CONTAMINANTS BY CONSIDERATION OF TURBULENT CHARACTERISTICS OF WSMR

By Frank V. Hansen



# ATMOSPHERIC SCIENCES LABORATORY

WHITE SANDS MISSILE RANGE, NEW MEXICO

CEEAVINGBEOGE

Distribution of this report is unlimited.

ECOM

UNITED STATES ARMY ELECTRONICS COMMAND

# PREDICTING DIFFUSION OF ATMOSPHERIC CONTAMINANTS BY CONSIDERATION OF TURBULENT CHARACTERISTICS OF WSMR

By

FRANK V. HANSEN

ECCM-5170

January 1968

DA TASK 1T014501B53A-10

ATMOSPHERIC SCIENCES LABORATORY
White Sands Missile Range, New Mexico

Distribution of this report is unlimited


### **ABSTRACT**

The relationships between the turbulent intensities about the mean flow in the lowest layers of the atmosphere at White Sands Missile Range, New Mexico, and climatological data are discussed in terms of predicting the diffusion of contaminants into the atmosphere. If the turbulent fluctuations observed in the lower portion of the boundary layer are considered representative, then correlations with climatological summaries may be used as a rudimentary estimator for estimating the diffusion power of the atmosphere. A simple empirical diffusion model for general use is also given.

# CONTENTS

	Page
ABSTRACT	iii
INTRODUCTION	1
PURPOSE OF DIFFUSION STUDY	2
THE SITE	2
TURBULENT CHARACTERISTICS OF WSMR	2
TERRAIN EFFECTS ON THE MEAN WIND	15
AN EMPIRICAL DIFFUSION PREDICTION MODEL	15
DISCUSSION AND SUMMARY	23
REFERENCES	24

Preceding Page Blank



V

#### INTRODUCTION

Diffusion of aerosols and particulate matter into the atmosphere from an instantaneous or continuous point source can be adequately predicted over short distances if restrictions are placed on the prediction model. Since problems of diffusion are of an individualistic nature, i.e., the conditions for diffusion vary from locale to locale, prediction schemes are normally determined empirically for a specific location and are based upon the theoretical models of Hay and Pasquill [1959] or Sutton [1932, 1934, 1947].

The development of a prediction scheme normally involves detailed studies of the area in question with respect to the meteorological and climatological characteristics and the use of tracer materials and detection systems to determine dispersion patterns. The actual field testing is normally easy to accomplish; however, the extraction of intelligence from the resulting data in terms of the meteorological situation are sometimes meaningless and obscure. The major causes of a poor data fit to theoretical considerations usually stem from the fact that the prediction models are based upon stationary conditions and homogeneous underlying terrain. Since most locations where there is a diffusion problem are less than ideal from a meteorological standpoint, the empirical solutions for atmospheric dispersion of gases or matter are necessarily based upon gross statistics and regression techniques and fail to account for the heterogeneous effects that lead to prediction errors that can over- or underestimate by a factor of two to four.

The scatter about a regression curve can be considered a measure of nonuniform effects upon the dispersion of contaminants into the atmosphere. The processes determining the heterogeneity of flow across a surface are not completely understood; consequently, it is difficult to compensate for this in terms of diffusion theory and its application. Thus, timely and realistic predictions of dispersion corridors and concentrations of contaminants at any point downstream of the source must necessarily be in the three-sigma range to allow for the effects of nonhomogeneous conditions.

A diffusion study for any locale, then, must include a micrometeorological study that considers all factors of the mean flow in steady and nonsteady state conditions and over a wide stability range. The vertical fluxes of heat, momentum, and matter into the atmosphere, and of course all the other turbulent parameters must be determined for the particular surface over which the diffusion will occur. Surface discontinuities that cause horizontal gradients and nonequilibrium profiles and fluxes great distances downstream must be compensated for in the analysis of test data.

#### PURPOSE OF DIFFUSION STUDY

The Atmospheric Sciences Laboratory, White Sands Missile Range (WSMR), New Mexico, is currently engaged in a study of the meteorological factors that affect the diffusion of toxic liquid propellants into the atmosphere. The purpose of the study is to provide the National Aeronautics and Space Administration White Sands Test Facility (WSTF) with timely and representative information concerning possible hazard corridors and concentration of toxic vapors in the vicinity of the Apollo site.

The purpose of this paper is to present some preliminary data concerning the characteristics of the planetary boundary layer that may be applied to diffusion into the atmosphere. Also included is an empirical diffusion model that may be used with some success over a rough, nonhomogeneous terrain.

#### THE SITE

White Sands Missile Range is located in South-Central New Mexico, almost wholly in the Tularosa Basin. The western boundaries of the range extend across the San Andres Range of the southern Rockies into the eastern portion of the Rio Grande Valley and the Jornada Del Muerto. The Apollo site is located on the western slopes of the San Andres and is mainly on the talus of the mountain range which is characterized by numerous arroyos, rocky soil, and sparse vegetation. The locale is heterogeneous from the micrometeorological standpoint. Man-made obstructions such as buildings and other modifications to the terrain contribute to the overall heterogeneity.

#### TURBULENT CHARACTERISTICS OF WSMR

The diffusive power of the atmosphere is directly related to the turbulent characteristics of the planetary boundary layer, and particularly to the vertical temperature gradient, the mean wind speed and the direction. The logical starting point, then, of any diffusion study is to review the turbulent characteristics of the atmosphere for the locale in question; keeping in mind of course, the differences that may exist owing to possible separation of the data collection site and the area of interest.

The investigation of the turbulent characteristics in the vicinity of WSMR has been quite productive in that large data samples have been collected and thoroughly analysed. A goodly portion of the data are applicable to the Apollo side of the mountains, even though they were collected in the Tularosa basin portion of WSMR.

It has been shown by Monin-Obukhov [1954], Panofsky [1961, 1963] Ellison [1957] and others that atmospheric turbulence obeys the laws of dynamic similarity of flows. In the surface boundary layer and to lesser extent in the planetary boundary layer proper, the wind and temperature profiles can be shown to be given by, respectively,

$$\frac{2}{L} = S \operatorname{Ri} K_{H} / K_{M} \tag{1}$$

and

$$\frac{z}{L} = R \operatorname{Ri} \left( K_{H} / K_{M} \right)^{2} \tag{2}$$

where z/L is the Monin-Obukhov scaling ratio, Ri is the gradient Richardson number, S is a dimensionless wind shear, R is a dimensionless lapse rate and  $K_H/K_M$  is the ratio of the exchange coefficients for heat and momentum. Batchelor [1953] has shown that the Richardson number is the basic parameter of similarity for the boundary layer, and it has been demonstrated by Hansen [1966], and Swanson and Cramer [1965] that the turbulent characteristics of the boundary layer are directly related to this basic stability parameter. The Richardson number is usually expressed as

$$Ri = \frac{g}{\theta} \frac{\partial \overline{\theta}/\partial z}{(\partial \overline{V}/\partial_z)^2}$$
 (3)

where g is the acceleration of gravity,  $\overline{\theta}$  is potential temperature,  $\overline{V}$  is the mean horizontal wind speed and z is height. From the similarity theory it may be shown that the Richardson number is also a function of height and its vertical distribution governed by

(Ri) = 
$$\frac{\sum \text{Ri}(z)}{\sum z}$$
 (j) (4)


where (Ri) denotes the height derivative of Ri. An investigation of the distribution of Ri with time as well as height by Hansen and Lang [1966] reveals that eq. (4) adequately describes the height distribution of Ri to the top of the surface boundary layer which, depending upon stability, is some 10 to 40 neters above the surface. This is shown in Figures 1 and 2 for both homogeneous and nonhomogeneous flow conditions. Figures 1 and 2 reveal that critical values of Ri exist, especially in the unstable regime. It has been found by Dyer [1965] and Hansen [1967] that critical Ri is in the range -0.70 > Ri > -1.2. In the stable regime, critical Ri is thought to lie between Ri = 0.20 and Ri = 0.40. Outside these limits are regimes of local nonstationarity known as the "undulant" regime for very stable conditions and the "windless convection" regime for the extremely unstable case.

It was also found that temporally the distribution of Ri with height was as depicted in Figure 3. It is seen that the Richardson number varies with the energy balance of the boundary layer, which is best shown in Figure 4. From Figures 3 and 4 it is seen that the atmosphere is almost wholly thermally stratified at all times and that neutral conditions exist only for very short periods near sunrise and sunset. It is interesting to note that the Richardson number at 3 and 6 meters is almost perfectly correlated with the incoming short wave and net radiation curves, indicating that the stability of the atmosphere at these heights reflects steady-state conditions much better than some other reference height, say 1.5 meters.

If the basic similarity parameter is taken to be the Richardson number, and if it is taken to be an independent variable in the boundary layer, eqs. (1) and (2) may be integrated to find the wind and temperature profiles. Substituting in and integrating eq. (1) yields

$$\nabla = \frac{u_*}{k} \left[ \ln \frac{z}{z_0} - \psi \left( z/L \right) \right]$$
 (5)

which is known as the log-linear wind profile, where u, is a stress term, k is Karman's constant, z is the roughness length and  $\psi(z/L)$  is a universal function. A similar expression may be written for the temperature profile, although not in such a universal form. The turbulent characteristics of the boundary layer may then be expressed as deviations about the mean profile as functions of Fi.



ABSOLUTE VALUE OF THE RICHARDSON NUMBER VS HEIGHT FOR EIGHT STABILITY CONDITIONS FOR HOMOGENEOUS FLOW.




FIGURE 2 ABSOLUTE VALUE OF THE RICHARDSON NUMBER VERSUS HEIGHT FOR SEVEN STABILITY CONDITIONS FOR NONHOMOGENEOUS FLOW.





FIG 3 THE RICHARDSON NUMBER AS A FUNCTION OF HEIGHT AND TIME



THE MCHARDSON NR. AT SELECTED WEIGHTS VERSUS NET AND INCOMING SHORT. WAVE RADIATION. F16.4

Swanson and Cramer [1965], Hansen [1966] and Hansen and Lang [1967] have shown that the turbulent wind and temperature fluctuations about the means are directly related to the stability regime. It was found that the orthogonal components of the horizontal wind were functions of stability and to a lesser extent functions of surface roughness. The turbulent intensity was also found to be height dependent in the form

$$Gx_* Gy = Cz^p$$
 (6)

where Gx and Gy are defined as

$$Gx = \sigma u^{\dagger} / \overline{V} \qquad (7)$$

$$Gy = \sigma v' / \overline{V} \stackrel{\sim}{\sim} \sigma A_{R}$$
 (8)

where ou' and ov' are the standard deviations of the longitudinal and lateral eddy velocities and  $\sigma A_R$  is the standard deviation about the mean wind direction in radians. The constant C is a function of surface roughness while the exponent p is related to the exponent of the conjugate power law wind profile and is a function of Ri.

llansen and Lang's [1967] results are shown in Figures 5 through 8. It will be noted that an inflection occurs in the  $\sigma A$  versus z regression of Figure 8 for the stable case. This is attributed to a surface roughness discontinuity upstream of the research tower as reported by llansen and llansen [1965]. The inflection in the curve is the result of excessive surface stress downstream of the discontinuity and the lack of vertical transport of momentum in stable flow conditions through the internal momentum boundary.

Turbulent temperature fluctuations over a nonhomogeneous surface are shown in Figure 9 as a function of the Richardson number. It is seen that in the region below the internal momentum boundary the quantity  $\sigma T/\partial\theta/\partial z$  does not exhibit the predicted characteristics as indicated by Tsvang [1960] and Lumley and Panofsky [1964]. Above the interface height of 8 to 10 meters, the scaling temperature  $T^*(T^*\chi \sigma T/\partial\theta/\partial z)$  has the predictable characteristic of being nearly unity in unstable flow and independent of height in stable conditions. The dashed portions of this curve indicate local nonstationarity




FIGURE 5 HEIGHT DEPENDENCE OF THE LONGITUDINAL INTENSITY OF TURBULENCE DURING UNSTABLE STRATIFICATION.




FIGURE 6 HEIGHT DEPENDENCE OF THE LONGITUDINAL INTENSITY OF TURBULENCE DURING STABLE STRATIFICATION




FIGURE 7 HEIGHT DEPENDENCE OF THE LATERAL INTENSITY OF TURBULENCE DURING UNSTABLE STRATIFICATION





FIGURE 8 HEIGHT DEPENDENCE OF THE LATERAL INTENSITY OF TURBULENCE DURING STABLE STRATIFICATION



effects upon the experimental data. On the stable side, it appears that undulance has the same effects as the nonequilbrium flow beneath the boundary.

#### TERRAIN EFFECTS ON THE MEAN WIND

The terrain of WSMR and particularly the proximity of a mountain range to the experimental site has a profound effect upon the annual wind regime at WSMR. Hansen and Lang [1966b, 1966c] found that the annual wind regime could be expressed in terms of three seasonal regimes. These consisted of a winter regime, a two-part opring regime and a combined summer-fall regime. The winter and spring seasons are characterized by generally strong westerly flow with the summer-fall regime being generally southeasterly. All seasons reflect a significant percentage of drainage winds from the mountain slopes during the nocturnal hours. The percent frequency of occurrence of wind direction for all seasons is shown in Figures 10 through 13.

Other terrain effects such as the roughness length have been investigated. Hansen [1967] has found that the roughness length of the terrain about the research tower varies with wind direction and with the season. The seasonal change in roughness is shown in Figure 14. It will be noted that the curve approximates the growing season in that the largest values of  $z_0$  are found in July and August when the vegetive canopy is most dense.

#### AN EMPIRICAL DIFFUSION PREDICTION MODEL

Experimental evidence shows that three meteorological parameters should be considered in characterizing the rate of small scale atmospheric diffusion:

- (1) The mean wind speed  $\overline{V}$  as an indicator of the downstream travel of the plume emitted from a continuous source.
- (2) The standard deviation of wind direction  $\sigma\Lambda$  as an indicator of the lateral rate of mixing.
- (3) The temperature gradient  $\Delta T$  as an indicator of the vertical rate of mixing.

Other quantities of interest are the peak or maximum concentration  $\mathbf{C}_{\mathrm{D}}$  at some distance x downstream of the source and the related source




FIGURE 10 PERCENT FREQUENCY OF OCCURRENCE OF WIND DIRECTION, WINTER




FIGURE II PERCENT FREQUENCY OF OCCURRENCE OF WIND DIRECTION, SPRING (PARTI)




FIGURE 12 PERCENT FREQUENCY OF OCCURRENCE OF WIND DIRECTION, SPRING (PART 2)





FIGURE 13 PERCENT FREQUENCY OF OCCURRENCE OF WIND DIRECTION, SUMMER-FALL



strength (). In the development of a regression equation for predicting diffusion, a normalized peak concentration  $C_p/Q$  at distance x is generally the goal.

The regression equation usually takes the form

$$C_{p}/Q = Kx^{a} \overline{V}^{b} \sigma A^{c} (\Delta T + t)^{d}$$
(9)

where K, a, b, c, d are parameters of fit and t is a constant added to  $\Delta T$  to avoid raising a negative number to a power.

Application of linear multiple regression techniques requires taking the logarithm of eq (9) and applying standard least-squares analysis to the following:

$$\log C_{p}/Q = \log K + a \log x + b \log \overline{V} + c \log \sigma A + d \log(\Delta T + t)$$
 (10)

since the normalized peak concentration could be linear with respect to the meteorological variables but not linear in distances.

A general purpose form of equation (9) for use in nonhomogeneous terrain has been developed by Miller and Miller [1963] in the form

$$C_p/Q = 0.000175 \times x^{-1.95} (\Delta T + 10)^{4.95}$$
 (11)

where  $C_p$  is the peak concentration in gms m<sup>-3</sup> at a height of about 1.5 meters above the surface at a downwind travel distance, x, in meters, Q is the source strength in gm sec<sup>-1</sup> and  $\Delta T$  is the temperature difference in deg. F between 16.5 and 1.8 meters above the ground. Eq. (11) was developed from experimental data simulating ground-level continuous point sources.

It will be noted that the mean wind speed V and the standard deviation of wind direction are not included in eq. (11). It was found by Haugen and Taylor [1963] and Miller and Miller [1963] that while V is correlated with  $C_p/Q_r$ , its contribution to the predictability after the temperature differential is included was not significant. The  $\sigma\Lambda$  term was dropped owing to the fact that  $\tau_0$  satisfactory

method of obtaining  $\sigma A$  from an analogue recorder was available. Estimates for  $\sigma A$  will be introduced below.

For direct application, eq (11) must be converted to units which may be used conveniently. The conversion is relatively simple for contaminants emitted as a ground-level continuous point source. In the case of a gaseous contaminant,  $C_p/Q$  should be known in terms of parts per million of the contaminant per pound per minute released into the atmosphere, a value that is dependent upon the diffusive power of the atmosphere and the physical characteristics of the contaminant. Under standard conditions, 1 ppm  $1b^{-1}$  min equals the gram molecular weight of the gas times  $5.89 \times 10^{-6}$  sec m<sup>-3</sup>. To obtain parts per million per pound per minute,  $C_p/Q$  in sec m<sup>-3</sup>, is multiplied by  $1.70 \times 10^5$  and divided by the gram molecular weight.

In terms of travel distance for normalized peak concentrations eq (11) may be restated as

$$x(in feet) = 2.62(C_p/Q)^{-0.513} [\Delta T+10]^{2.53}$$
 (12)

and

$$x(in feet) = 2.20 (C_p/Q)^{-0.513} [\Delta T+10]^{2.53}$$
 (13)

Eqs. (12) and (13) are valid for two well-known propellants, nitrogen tetroxide and unsymmetrical dimethylhydrazine. For other fuels or oxidizers, the constants of eqs. (12) and (13) must be modified according to the scheme outlined above.

Travel time of the contaminant to a particular location can be determined from the mean wind speed. The width of the hazard corridor can be estimated from a wind direction trace by

$$W = C R \tag{14}$$

where W is the width of the cerridor, R is the range of the wind direction, and C is an empirical constant determined for a locale and a particular recorder speed. Eq. (14) yields equivalent results obtained by Haugen and Taylor [1963] who showed that the width of the hazard corridor could be estimated from W=8 oA with 95% confidence that this value would not be exceeded.

#### DISCUSSION AND SUMMARY

If the turbulent characteristics of the boundary layer as observed at WSMR can be considered typical of an area that includes the WSTF (Apollo) locale, some insight into the diffusion of a contaminant into the atmosphere is provided. If the WSMR turbulence intensities are typical, then climatological type data at WSTF may be used to investigate the diffusive power of the atmosphere, alleviating any need for elaborate multilevel tower systems which are a necessity at WSMR.

The only exception to this approach would be the drainage wind pattern during the nocturnal hours. It is clear that the drainage pattern at WSTF would not be the same as it is at WSMR. This is because WSMR and WSTF are on opposite slopes of the mountain range. It appears likely that the nocturnal drainage on the western slopes of the San Andres would be almost a mirror image of the eastern slope drainage pattern with few exceptions.

The use of the Miller and Miller [1963] diffusion equation for predicting concentrations and hazard corridors for WSTF is perhaps not completely justified but is a useful tool for estimating the dispersion of contaminants in the atmosphere. The development of a more precise empirical model for WSTF would entail a very complex and costly experiment.

This analysis indicates that the turbulent characteristics of the atmosphere as observed at White Sands are representative of larger areas. Under these conditions then, the diffusive power of the atmosphere at the Apollo site may be estimated from rudimentary climatological data used in conjunction with the WSMR data.

#### REFERENCES

- Batchelor, G. K. [1953] The conditions for dynamic similarity of motion of a frictionless perfect-gas atmosphere. J. Roy Meteor. Soc. 79, 224.
- Dyer, A. J. [1965] The flux-gradient relation for turbulent heat transfer in the lower atmosphere. J. Roy. Meteor. Soc. 91, 151.
- Ellison, T. H. [1957] Turbulent transport of heat and momentum from an infinite rough plane. J. Fl. Mech. 2, 456.
- Hansen, F. V. [1966] The lateral intensity of turbulence as a function of stability. ECOM 5093, Atmospheric Sciences Laboratory, White Sands Missile Range, New Mexico.
- Ilansen, F. V. [1967] The aerodynamic roughness of the complex terrain of White Sands Missile Range, New Mexico. Atmospheric Sciences Laboratory, White Sands Missile Range, New Mexico.
- Hansen, F. V. [1967] Spacial and temporal distribution of the gradient Richardson number in the surface and planetary layers. ECOM-5123, Atmospheric Sciences Laboratory, White Sands Missile Range, New Mexico.
- Hansen, F. V. and P. S. Hansen [1965] The formation of an internal boundary over heterogeneous terrain. Atmospheric Sciences Laboratory, White Sands Missile Range, New Mexico.
- Hansen, F. V. and V. D. Lang [1966a] The Richardson number as a function of time and vertical distribution. Presented at the AGU Western National Meeting, Sept 6-9,1966, Los Angeles, California
- Hansen, F. V. and V. D. Lang [1966b] The wind regime in the first 62 meters of the atmosphere. ECOM 5058. Atmospheric Sciences Laboratory, White Sands Missile Range, New Mexico.
- Hansen, F. V. and V. D. Lang [1966c] Atmospheric turbulence over a nonhomogeneous terrain in a semiarid region. Presented at the 47th annual meeting, Pacific Division, AAAS, 13-18 June 1966, Seattle, Washington.
- Hansen, F. V. and V. D. Lang [1967] The turbulent intensity characteristics of the first 62 meters of the boundary layer at White Sands Missile Range. Presented at the 47th annual meeting, AMS. 23-26 January 1967, New York.

- Haugen, D. A. and J. H. Taylor [1963] The Ocean Breeze and Dry Gulch Diffusion programs, II. Rpt 63-791 (II) AFCRL, Bedford, Mass.
- Hay, J. S. and F. Pasquill [1959] Diffusion from a continuous source in relation to the spectrum and scale of turbulence. Ad Geophys 6 Academic Press, New York.
- Lumley, J. H. and H. A. Panofsky [1964] The Structure of Atmospheric Turbulence, John Wiley & Sons, New York.
- Miller, R. L. and F. H. Miller [1963] Meteorological support to Titan II propellant transfer operations. AFCRL, Bedford, Mass.
- Morin, A. S. and A. M. Obukhov [1954] Basic regularity in turbulent mixing in the surface layer of the atmosphere. <u>Trudy Geophys.</u> Inst. ANSSR, 24, 163.
- Panofsky, H. A. [1961] An alternative derivation of the diabatic wind profile. J. Roy. Meteor. Soc. 87, 109.
- Panofsky, II. A. [1963] Determination of stress from wind and temperature measurements. J. Roy. Meteor. Soc. 89, 85.
- Swanson, R. N. and H. E. Cramer [1965] A study of lateral and longitudinal intensities of turbulence. J. Appl. Meteor. 4, 409.
- Sutton, O. G. [1932] A theory of eddy diffusion in the atmosphere. Proc. Roy. Soc. A, 135, 143.
- Sutton, O. G. [1934] Wind structure and evaporation in a turbulent atmosphere. Proc. Roy. Soc. A, 146, 701.
- Sutton, O. G. [1947] The theoretical distribution of airhorne pollution from factory chimneys. Q. J. Roy. Meteor. Soc. 73, 426.
- Tsvang, L. R. [1960] Measurements of temperature pulse frequency spectra in the surface layer of the atmosphere. <u>Izvestia</u> ANSSSR, Geophys. Ser. 1960, 8, 1252.

## ATMOSPHERIC SCIENCES RESEARCH PAPERS

- Webb, W.L., "Development of Droplet Size Distributions in the Atmosphere." June 1954.
- Hansen, F. V., and H. Rachele, "Wind Structure Analysis and Forecasting Methods for Rockets," June 1954.
- Webb, W. L., "Net Electrification of Water Droplets at the Earth's Surface," J. Meteorol., December 1954.
- Mitchell, R., "The Determination of Non-Ballistic Projectile Trajectories," March 1955.
- Webb, W. L., and A. McPike, "Sound Ranging Technique for Determining the Trajectory of Supersonic Missiles," ±1, March 1955.

  Mitchell, R., and W. L. Webb, "Electromagnetic Radiation through the Atmosphere," ±1, April 1955. 5.
- Webb, W. L., A. McPike, and H. Thompson, "Sound Ranging Technique for Determining the Trajectory of Supersonic Missiles," #2, July 1955.
- Barichivich, A., "Meteorological Effects on the Refractive Index and Curvature of Microwaves in the Atmosphere," August 1955.

  Webb, W. L., A. McPike and H. Thompson, "Sound Ranging Technique for Deter-8.
- mining the Trajectory of Supersonic Missiles," #3, September 1955.
- Mitchell, R., "Notes on the Theory of Longitudinal Wave Motion in the Atmosphere," February 1956.
- Webb, W. L., "Particulate Counts in Natural Clouds," J. Meteorol., April 1956. Webb, W. L., "Wind Effect on the Aerobee," = 1, May 1956. 11.
- 12.
- 13.
- Rachele, H., and L. Anderson, "Wind Effect on the Aerobee," #2, August 1956. Beyers, N., "Electromagnetic Radiation through the Atmosphere," #2, January 1957. Hansen, F. V., "Wind Effect on the Aerobee," #3, January 1957. 14.
- 15.
- 16.
- Kershner, J., and H. Bear, "Wind Effect on the Aerobee," #4, January 1957. Hoidale, G., "Electromagnetic Radiation through the Atmosphere," #3, February 17. 1957.
- Querfeld, C. W., "The Index of Refraction of the Atmosphere for 2.2 Micron Radi-18. ation," March 1957.
- 19. White, Lloyd, "Wind Effect on the Aerobee," =5, March 1957.
- Kershner, J. G., "Development of a Method for Forecasting Component Ballistic 20. Wind," August 1957.
- Layton, Ivan, "Atmospheric Particle Size Distribution," December 1957. 21.
- Rachele, Henry and W. H. Hatch, "Wind Effect on the Aerobee," #6, February 1958.
- Beyers, N. J., "Electromagnetic Radiation through the Atmosphere," #4, March 23. 1958.
- Prosser, Shirley J., "Electromagnetic Radiation through the Atmosphere," #5, 24. April 1958.
- 25. Armendariz, M., and P. H. Taft, "Double Theodolite Ballistic Wind Computations," June 1958.
- Jenkins, K. R. and W. L. Webb, "Rocket Wind Measurements," June 1958. Jenkins, K. R., "Measurement of High Altitude Winds with Loki," July 1958. 27.
- Hoidale, G., "Electromagnetic Propagation through the Atmosphere," #6, February 1959.
- 29. McLardie, M., R. Helvey, and L. Traylor, "Low-Level Wind Profile Prediction Techniques," ±1. June 1959.

  Lamberth, Roy, "Gustiness at White Sands Missile Range," ±1, May 1959.
- 30.
- Beyers, N. J., B. Hinds, and G. Hoidale, "Electromagnetic Propagation through the Atmosphere," #7, June 1959.
- 32. Beyers, N. J., "Radar Refraction at Low Elevation Angles (U)," Proceedings of the
- Army Science Conference, June 1959. White, L., O. W. Thiele and P. H. Taft, "Summary of Ballistic and Meteorological Support During IGY Operations at Fort Churchill, Canada," August
- 'Hainline, D. A., "Drag Cord-Aerovane Equation Analysis for Computer Application," August 1959.
- Hoidale, G. B., "Slope-Valley Wind at WSMR," October 1959.
- Webb, W. L., and K. R. Jenkins, "High Altitude Wind Measurements," J. Meteorol., 16, 5, October 1959.

- White, Lloyd, "Wind Effect on the Aerobee," #9, October 1959.
- Webb, W. L., J. W. Coffman, and G. Q. Clark, "A High Altitude Acoustic Sensing System," December 1959.
- Webb, W. L., and K. R. Jenkins, "Application of Meteorological Rocket Systems," J. Geophys. Res., 64, 11, November 1959. 39.
- Duncan, Louis, "Wind Effect on the Aerobee," #10, February 1960. 40.
- Helvey, R. A., "Low-Level Wind Profile Prediction Techniques," #2, February 1960. 41.
- Webb, W. L., and K. R. Jenkins, "Rocket Sounding of High Altitude Parameters,"

  Proc. GM Rel. Symp., Dept. of Defense, February 1960. 42.
- Armendariz, M., and H. H. Monahan, "A Comparison Between the Double Theodolite and Single-Theodolite Wind Measuring Systems," April 1960.
- 44.
- Jenkins, K. R., and P. H. Taft, "Weather Elements in the Tularosa Basin," July 1960. Beyers, N. J., "Preliminary Radar Performance Data on Passive Rocket-Borne Wind Sensors," IRE TRANS, MIL ELECT, MIL-4, 2-3, April-July 1960. Webb, W. L., and K. R. Jenkins, "Speed of Sound in the Stratosphere," June 1960.
- 46.
- Webb, W. L., K. R. Jenkins, and G. Q. Clark, "Rocket Sounding of High Atmosphere Meteorological Parameters," IRE Trans. Mil. Elect., MIL-4, 2-3, April-July 1960.
- Helvey, R. A., "Low-Level Wind Profile Prediction Techniques," #3, September 48.
- Beyers, N. J., and O. W. Thiele, "Meteorological Wind Sensors," August 1960. 49.
- Armijo, Larry, "Determination of Trajectories Using Range Data from Three Non-50. colinear Radar Stations," September 1960.
- Carnes, Patsy Sue, "Temperature Variations in the First 200 Feet of the Atmo-51. sphere in an Arid Region," July 1961.

  Springer, H. S., and R. O. Olsen, "Launch Noise Distribution of Nike-Zeus Mis-
- 52. siles," July 1961.
  Thiele, O. W., "Density and Pressure Profiles Derived from Meteorological Rocket
- 53. Measurements," September 1961.
- Diamond, M. and A. B. Gray, "Accuracy of Missile Sound Ranging," November 54. 1961.
- Lamberth, R. L. and D. R. Veith, "Variability of Surface Wind in Short Distances." 55. #1, October 1961
- Swanson, R. N., "Low-Level Wind Measurements for Ballistic Missile Application," 56. January 1962.
- Lamberth, R. L. and J. H. Grace, "Gustiness at White Sands Missile Range," #2. January 1962.
- Swanson, R. N. and M. M. Hoidale, "Low-Level Wind Profile Prediction Techniques," #4, January 1962. 58.
- Rachele, Henry, "Surface Wind Model for Unguided Rockets Using Spectrum and 59. Cross Spectrum Techniques," January 1962.
- Rachele, Henry, "Sound Propagation through a Windy Atmosphere." #2. Febru-60. ary 1962.
- Webb, W. L., and K. R. Jenkins, "Sonic Structure of the Mesosphere," J. Acous. 61.
- Soc. Amer., 34, 2, February 1962.

  Tourin, M. H. and M. M. Hoidale, "Low-Level Turbulence Characteristics at White Sands Missile Range," April 1962. 62.
- Miers, Bruce T., "Mesospheric Wind Reversal over White Sands Missile Range." March 1962.
- Fisher, E., R. Lee and H. Rachele, "Meteorological Effects on an Acoustic Wave within a Sound Ranging Array," May 1962. 64.
- **6**5.
- Walter, E. L., "Six Variable Ballistic Model for a Rocket," June 1962. Webb, W. L., "Detailed Acoustic Structure Above the Tropopause," J. Applied Meteorol., 1, 2, June 1962.
- Jenkins, K. R., "Empirical Comparisons of Meteorological Rocket Wind Sensors," J. 67. Appl. Meteor., June 1962.

  Lamberth, Roy, "Wind Variability Estimates as a Function of Sampling Interval,"
- 68. July 1962.
- Rachele, Henry, "Surface Wind Sampling Periods for Unguided Rocket Impact Pre-69. diction," July 1962.
- Traylor, Larry, "Coriolis Effects on the Aerobee-Hi Sounding Rocket," August 1962.
- 71. McCoy, J., and G. Q. Clark, "Meteorological Rocket Thermometry," August 1962. 72. Rachele, Henry, "Real-Time Prelaunch Impact Prediction System," August 1962.

- Beyers, N. J., O. W. Thiele, and N. K. Wagner, "Performance Characteristics Meteorlogical Rocket Wind and Temperature Sensors," October 1962.
- Coffman, J., and R. Price, "Some Errors Associated with Acoustical Wind Measurements through a Layer," October 1962.
- Armendariz, M., E. Fisher, and J. Serna, "Wind Shear in the Jet Stream at WS-MR," November 1962.
- Armendariz, M., F. Hansen, and S. Carnes, "Wind Variability and its Effect on Rocket Impact Prediction," January 1963.
- Querfeld, C., and Wayne Yunker, "Pure Rotational Spectrum of Water Vapor, I: Table of Line Parameters," February 1963.
- Webb, W. L., "Acoustic Component of Turbulence," J. Applied Meteorol., 2, 2, 78. April 1963.
- 79. Beyers, N. and L. Engberg, "Seasonal Variability in the Upper Atmosphere," May 1963.
- 80. Williamson, L. E., "Atmospheric Acoustic Structure of the Sub-polar Fall," May 1963.
- Lamberth, Roy and D. Veith, "Upper Wind Correlations in Southwestern United 81. States," June 1963.
- Sandlin, E., "An analysis of Wind Shear Differences as Measured by AN/FPS-16 82.
- Radar and AN/GMD-1B Rawinsonde," August 1963.

  Diamond, M. and R. P. Lee, "Statistical Data on Atmospheric Design Properties Above 30 km," August 1963. 83.
- Thiele, O. W., "Mesospheric Density Variability Based on Recent Meteorological 84.
- Rocket Measurements," J. Applied Meteorol., 2, 5, October 1963.

  Diamond, M., and O. Essenwanger, "Statistical Data on Atmospheric Design Properties to 30 km," Astro. Aero. Engr., December 1963.

  Hansen, F. V., "Turbulence Characteristics of the First 62 Meters of the Atmo-85.
- 86. sphere," December 1963.
- Morris, J. E., and B. T. Miers, "Circulation Disturbances Between 25 and 70 kilo-87. meters Associated with the Sudden Warming of 1963," J. of Geophys. Res., January 1964.
- Thiele, O. W., "Some Observed Short Term and Diurnal Variations of Stratospheric Density Above 30 km." January 1964. 88.
- Sandlin, R. E., Jr. and E. Armijo, "An Analysis of AN/FPS-16 Radar and AN/ 89. GMD-1B Rawinsonde Data Differences," January 1964.
- Miers, B. T., and N. J. Beyers, "Rocketsonde Wind and Temperature Measurements Between 30 and 70 km for Selected Stations," J. Applied Mete-90. orol., February 1964.
- Webb, W. L., "The Dynamic Stratesphere," Astronautics and Aerospace Engineer-91. ing, March 1964.
- Low, R. D. H., "Acoustic Measurements of Wind through a Layer," March 1964. Diamond. M.. "Cross Wind Effect on Sound Propagation," J. Applied Meteorol., 93. April 1964.
- Lee, R. P., "Acoustic Ray Tracing," April 1964. 94.
- Reynolds, R. D., "Investigation of the Effect of Lapse Rate on Balloon Ascent Rate." 95. May 1964.
- Webb, W. L., "Scale of Stratospheric Detail Structure," Space Research V, May 96. 1964.
- Barber, T. L., "Proposed X-Ray-Infrared Method for Identification of Atmospheric Mineral Dust," June 1964.

  Thiele, O. W., "Ballistic Procedures for Unguided Rocket Studies of Nuclear Environ-97.
- 98. ments (U)," Proceedings of the Army Science Conference, June 1964. Horn, J. D., and E. J. Trawle, "Orographic Effects on Wind Variability," July 1964. 99.
- Hoidale, G., C. Querfeld, T. Hali, and R. Mireles, "Spectral Transmissivity of the Earth's Atmosphere in the 250 to 500 Wave Number Interval," #1, 100. September 1964.
- 101. Duncan, L. D., R. Ensey, and B. Engebos, "Athena Launch Angle Determination,"
- September 1964.
  Thiele, O. W., "Feasibility Experiment for Measuring Atmospheric Density Through 102. the Altitude Range of 60 to 100 KM Over White Sands Missile Range,' October 1964.
- Duncan, L. D., and R. Ensey, "Six-Degree-of-Freedom Digital Simulation Model for Unguided, Fin-Stabilized Rockets," November 1964.

Hoidale, G., C. Querfeld, T. Hall, and R. Mireles, "Spectral Transmissivity of the 104 Earth's Atmosphere in the 250 to 500 Wave Number Interval, November 1964.

Webb, W. L., "Stratospheric Solar Response," J. Atmos. Sci., November 1964. 105.

McCoy, J. and G. Clark, "Rocketsonde Measurement of Stratospheric Temperature," 106. December 1964.

- Farone, W. A., "Electromagnetic Scattering from Radially Inhomogeneous Spheres 107. as Applied to the Problem of Clear Atmosphere Radar Echoes," December 1964.
- 108. Farone, W. A., "The Effect of the Solid Angle of Illumination or Observation on the Color Spectra of 'White Light' Scattered by Cylinders," January 1965.

Williamson, L. E., "Seasonal and Regional Characteristics of Acoustic Atmospheres," 109.

J. Geophys. Res., January 1965.

Armendariz, M., "Ballistic Wind Variability at Green River, Utah," January 1965.

Low, R. D. H., "Sound Speed Variability Due to Atmospheric Composition," Janu-110. 111.

ary 1965. Querfeld, C. W., 'Mie Atmospheric Optics," J. Opt. Soc. Amer., January 1965. 112.

Coffman, J., "A Measurement of the Effect of Atmospheric Turbulence on the Coherent Properties of a Sound Wave," January 1965. 113.

Rachele, H., and D. Veith, "Surface Wind Sampling for Unguided Rocket Impact 114.

Prediction," January 1965.
Ballard, H., and M. Izquierdo, "Reduction of Microphone Wind Noise by the Gen-115. eration of a Proper Turbulent Flow," February 1965.

116.

Mireles, R., "An Algorithm for Computing Half Widths of Overlapping Lines on Experimental Spectra," February 1965.

Richart, H., "Inaccuracies of the Single-Theodolite Wind Measuring System in Ballistic Application," February 1965.

D'Arry M. "Theorytical and Practical Study.

D'Arcy, M., "Theoretical and Practical Study of Aerobee-150 Ballistics," March 118. 1965.

"Improved Method for the Reduction of Rocketsonde Temperature Da-119. McCoy, J., ta," March 1965. "Uniqueness Theorem in Inverse Electromagnetic Cylindrical Scatter-120. Mireles, R.,

ing," April 1965. Coffman, J., "The Focusing of Sound Propagating Vertically in a Horizontally Stra-

tified Medium," April 1965.

Farone, W. A., and C. Querfeld, "Electromagnetic Scattering from an Infinite Circular Cylinder at Oblique Incidence," April 1965.

Rachele, H., "Sound Propagation through a Windy Atmosphere," April 1965.

Miers, B., "Upper Stratospheric Circulation over Ascension Island," April 1965. 122.

123.

124. 125.

Rider, L., and M. Armendariz, "A Comparison of Pibal and Tower Wind Measurements," April 1965.

Hoidale, G. B., "Meteorological Conditions Allowing a Rare Observation of 24 Mi-126.

cron Solar Radiation Near Sea Level," Meteorol. Magazine, May 1965. Beyers, N. J., and B. T. Miers, "Diurnal Temperature Change in the Atmosphere 127. Between 30 and 60 km over White Sands Missile Range," J. Atmos. Sci., May 1965.

Querfeld, C., and W. A. Farone, "Tables of the Mie Forward Lobe," May 1965. 128.

Farone, W. A., Generalization of Rayleigh-Gans Scattering from Radially Inhomogeneous Spheres," J. Opt. Soc. Amer., June 1965. 129.

Diamond, M., "Note on Mesospheric Winds Above White Sands Missile Range," J. 130. Applied Meteorol., June 1965.

131.

Clark, G. Q., and J. G. McCoy, "Measurement of Stratospheric Temperature," J.

Applied Meteorol., June 1965.

Hall, T., G. Hoidale, R. Mireles, and C. Querfeld, "Spectral Transmissivity of the
Earth's Atmosphere in the 250 to 500 Wave Number Interval," #3, 132. July 1965.

McCoy, J., and C. Tate, "The Delta-T Meteorological Rocket Payload," June 1964. 133.

Horn, J. D., "Obstacle Influence in a Wind Tunnel," July 1965.

"An AC Probe for the Measurement of Electron Density and Collision 135. McCoy, J.,

Frequency in the Lower Ionosphere," July 1965.
Miers, B. T., M. D. Kays, O. W. Thiele and E. M. Newby, "Investigation of Short Term Variations of Several Atmospheric Parameters Above 30 KM," July 1965.

137. Serna, J., "An Acoustic Ray Tracing Method for Digital Computation," September 1965.

Webb, W. L., "Morphology of Noctilucent Clouds," J. Geophys. Res., 70, 18, 4463-138. 4475, September 1965.

Kays, M., and R. A. Craig, "On the Order of Magnitude of Large-Scale Vertical Motions in the Upper Stratosphere," J. Geophys. Res., 70, 18, 4453-4462, 139. September 1965.

140.

Rider, L., "Low-Level Jet at White Sands Missile Range," September 1965.

Lamberth, R. L., R. Reynolds, and Morton Wurtele, "The Mountain Lee Wave at White Sands Missile Range," Bull. Amer. Meteorol. Soc., 46, 10, Octo-141. ber 1965.

142. Reynolds, R. and R. L. Lamberth, "Ambient Temperature Measurements from Ra-

diosondes Flown on Constant-Level Balloons," October 1965.

McCluney, E., "Theoretical Trajectory Performance of the Five-Inch Gun Probe 143. System," October 1965.

Pena, R. and M. Diamond, "Atmospheric Sound Propagation near the Earth's Sur-144.

face," October 1965.

Mason, J. B., "A Study of the Feasibility of Using Radar Chaff For Stratospheric Temperature Measurements," November 1965. 145.

Diamond, M., and R. P. Lee, "Long-Range Atmospheric Sound Propagation," J. Geophys. Res., 70, 22, November 1965. 146.

Lamberth, R. L., "On the Measurement of Dust Devil Parameters," November 1965. 147.

Hansen, F. V., and P. S. Hansen, "Formation of an Internal Boundary over Heter-148. ogeneous Terralla," November 1965.

Webb, W. L., "Mechanics of Stratospheric Scasonal Reversals," November 1965.

U. S. Army Electronics R & D Activity, "U. S. Army Participation in the Meteoro-

150. logical Rocket Network," January 1966.

Rider, L. J., and M. Armendariz, "Low-Level Jet Winds at Green River, Utah," Feb-151. ruary 1966.

152. 153.

Webb, W. L., "Diurnal Variations in the Stratospheric Circulation," February 1966. Beyers, N. J., B. T. Miers, and R. J. Reed, "Diurnal Tidal Motions near the Stratopause During 48 Hours at WSMR," February 1966.

Webb, W. L., "The Stratospheric Tidal Jet," February 1966. 154.

Hall, J. T., "Focal Properties of a Plane Grating in a Convergent Beam," February 155. 1966.

Duncan, L. D., and Henry Rachele, "Real-Time Meteorological System for Firing of Unguided Rockets," February 1966. 156.

157. Kays, M. D., "A Note on the Comparison of Rocket and Estimated Geostrophic Winds

at the 10-mb Level," J. Appl. Meteor., February 1966.

Rider, L., and M. Armendariz, "A Comparison of Pibal and Tower Wind Measurements," J. Appl. Meteor., 5, February 1966. 158. 159.

Duncan, L. D., "Coordinate Transformations in Trajectory Simulations," February 1966.

Williamson, L. E., "Gun-Launched Vertical Probes at White Sands Missile Range," 160. February 1966. 161. Randhawa, J. S., Ozone Measurements with Rocket-Borne Ozonesondes," March

1966. 162.

Armendariz, Manuel, and Laurence J. Rider, "Wind Shear for Small Thickness Layers," March 1966.

Low, R. D. H., "Continuous Determination of the Average Sound Velocity over an Arbitrary Path," March 1966. 163.

Hansen, Frank V., "Richardson Number Tables for the Surface Boundary Layer," 164.

March 1966. 165.

Cochran, V. C., E. M. D'Arcy, and Florencio Ramirez, "Digital Computer Program for Five-Degree-of-Freedom Trajectory," March 1966.

Thiele, O. W., and N. J. Beyers, "Comparison of Rocketsonde and Radiosonde Temp-166.

eratures and a Verification of Computed Rocket onde Pressure and Den-

167.

eratures and a verification of Computed Rockettonge Pressure and Density," April 1966.

Thiele, O. W., "Observed Diurnal Oscillations of Pressure and Density in the Upper Stratosphere and Lower Mesosphere," April 1966.

Kays, M. D., and R. A. Craig, "On the Order of Magnitude of Large-Scale Vertical Motions in the Upper Stratosphere," J. Geophy. Res., April 1966.

Hansen, F. V., "The Richardson Number in the Planetary Boundary Layer," May 168.

169. 1966.

Ballard, H. N., "The Measurement of Temperature in the Stratosphere and Mesosphere," June 1966.

Hansen, Frank V., "The Ratio of the Exchange Coefficients for Heat and Momentum 170.

in a Homogeneous, Thermally Stratified Atmosphere," June 1966.

Hansen, Frank V., "Comparison of Nine Profile Models for the Diabatic Boundary 172. Layer," June 1966.

Rachele, Henry, "A Sound-Ranging Technique for Locating Supersonic Missiles," 173. May 1966

Farone, W. A., and C. W. Querfeld, "Electromagnetic Scattering from Inhomogeneous Infinite Cylinders at Oblique Incidence," J. Opt. Soc. Amer. 56, 4, 476-

480, April 1966.

Mireles, Ramon, "Determination of Parameters in Absorption Spectra by Numerical Minimization Techniques," J. Opt. Soc. Amer. 56, 5, 644-647, May 1966.

Reynolds, R., and R. L. Lamberth, "Ambient Temperature Measurements from Ra-

176. diosondes Flown on Constant-Level Balloons," J. Appl. Meteorol., 5, 3, 304-307, June 1966.

Hall, James T., "Focal Properties of a Plane Grating in a Convergent Beam," Appl.

Opt., 5, 1051, June 1966

Rider, Laurence J., "Low-Level Jet at White Sands Missile Range," J. Appl. Meteorol., 5, 3, 283-287, June 1966.

McCluney, Eugene, "Projectile Dispersion as Caused by Barrel Displacement in the 5-Inch Gun Probe System," July 1966. 178.

179.

Armendariz, Manuel, and Laurence J. Rider, "Wind Shear Calculations for Small Shear Layers," June 1966. 180.

Lamberth, Roy L., and Manuel Armendariz, "Upper Wind Correlations in the Cen-181. tral Rocky Mountains," June 1966.

Hansen, Frank V., and Virgil D. Lang, "The Wind Regime in the First 62 Meters of

182. the Atmosphere," June 1966.

183.

Randhawa, Jagir S., "Rocket-Borne Ozonesonde," July 1966.
Rachele, Henry, and L. D. Duncan, "The Desirability of Using a Fast Sampling Rate for Computing Wind Velocity from Pilot-Balloon Data," July 1966.
Hinds, B. D., and R. G. Pappas, "A Comparison of Three Methods for the Cor-184.

185.

rection of Radar Elevation Angle Refraction Errors," August 1966. Riedmuller, G. F., and T. L. Barber, "A Mineral Transition in Atmospheric Dust Transport," August 1966. 186.

Hall, J. T., C. W. Querfeld, and G. B. Hoidale, "Spectral Transmissivity of the Earth's Atmosphere in the 250 to 500 Wave Number Interval," Part 187.

IV (Final), July 1966.

Duncan, L. D. and B. F. Engebos, "Techniques for Computing Launcher Settings for Unguided Rockets," September 1966.

Duncan, L. D., "Basic Considerations in the Development of an Unguided Rocket 188.

189. Trajectory Simulation Model," September 1966.

Miller, Walter B., "Consideration of Some Problems in Curve Fitting," September

190.

191.

192.

193.

Cermak, J. E., and J. D. Horn, "The Tower Shadow Effect," August 1966.
Webb, W. L., "Stratospheric Circulation Response to a Solar Eclipse," October 1966.
Kennedy, Bruce, "Muzzle Velocity Measurement," October 1966.
Traylor, Larry E., "A Refinement Technique for Unguided Rocket Drag Coefficients," October 1966
Nusbaum, Henry, "A Reagent for the Simultaneous Microscope Determination of 194.

195.

Quartz and Halides," October 1966.

Kays, Marvin and R. O. Olsen, "Improved Rocketsonde Parachute-derived Wind Profiles," October 1966.

Engebos, Bernard F. and Duncan, Louis D., "A Nomogram for Field Determina-196.

197. tion of Launcher Angles for Unguided Rockets," October 1966.

Webb, W. L., "Midlatitude Clouds in the Upper Atmosphere," November 1966. 198. Hansen, Frank V., "The Lateral Intensity of Turbulence as a Function of Stability," 199. November 1966.

Rider, L. J. and M. Armendariz, "Differences of Tower and Pibal Wind Profiles," 200. November 1966.

201. Lee, Robert P., "A Comparison of Eight Mathematical Models for Atmospheric

Acoustical Ray Tracing," November 1966.

Low, R. D. H., et al., "Acoustical and Meteorological Data Report SOTRAN I and II," November 1966.

Hunt, J. A. and J. D. Horn, "Drag Plate Balance," December 1966. 203.

Armendariz, M., and H. Rachele, "Determination of a Representative Wind Profile from Balloon Data," December 1966. 204.

Hansen, Frank V., "The Aerodynamic Roughness of the Complex Terrain of White 205. Sands Missile Range," January 1967.

Morris, James E., "Wind Measurements in the Subpolar Mesopause Region," Jan-206. uary 1967.

Hall, James T., "Attenuation of Millimeter Wavelength Radiation by Gaseous 207. Water," January 1967.

Thiele, O. W., and N. J. Beyers, "Upper Atmosphere Pressure Measurements With 208. Thermal Conductivity Gauges," January 1967.

Armendariz, M., and H. Rachele, "Determination of a Representative Wind Profile from Balloon Data," January 1967. 209.

Hansen, F. V., "The Aerodynamic Roughness of the Complex Terrain of White Sands 210. Missile Range, New Mexico," January 1967.

D'Arcy, Edward M., "Some Applications of Wind to Unguided Rocket Impact Prediction," March 1967. 211.

Kennedy, Bruce, "Operation Manual for Stratosphere Temperature Sonde," March 212. 1967.

213. Hoidale, G. B., S. M. Smith, A. J. Blanco, and T. L. Barber, "A Study of Atmospheric Dust," March 1967.

Longyear, J. Q., "An Algorithm for Obtaining Solutions to Laplace's Titad Equa-214. tions," March 1967.

Rider, L. J., "A Comparison of Pibal with Raob and Rawin Wind Measurements," 215. April 1967. 216.

Breeland, A. H., and R. S. Bonner, "Results of Tests Involving Hemispherical Wind Screens in the Reduction of Wind Noise," April 1967. Webb. Willis L., and Max C. Bolen, "The D-region Fair-Weather Electric Field," 217.

April 1967. Kubinski, Stanley F., "A Comparative Evaluation of the Automatic Tracking Pilot-218.

Balloon Wind Measuring System," April 1967. 219.

Miller, Walter B., and Henry Rachele, "On Nonparametric Testing of the Nature of Certain Time Series," April 1967.

Hansen, Frank V., "Spacial and Temporal Distribution of the Gradient Richardson 220.

Number in the Surface and Planetary Layers," May 1967. 221.

Randhawa, Jagir S., "Diurnal Variation of Ozone at High Altitudes," May 1967. Ballard, Harold N., "A Review of Seven Papers Concerning the Measurement of 222. Temperature in the Stratosphere and Mesosphere," May 1967.

Williams, Ben H., "Synoptic Analyses of the Upper Stratospheric Circulation Dur-223. ing the Late Winter Storm Period of 1966," May 1967.

Horn, J. D., and J. A. Hunt, "System Design for the Atmospheric Sciences Office 224. Wind Research Facility," May 1967. Miller, Walter B., and Henry Rachele, "Dynamic Evaluation of Radar and Photo 225.

Tracking Systems, " May 1967. 226.

Bonner, Robert S., and Ralph H. Rohwer, "Acoustical and Meteorological Data Report - SOTRAN III and IV," May 1967. 227.

Rider, L. J., "On Time Variability of Wind at White Sands Missile Range, New Mexico," June 1967. 228.

Randhawa, Jagir S., "Mesospheric Ozone Measurements During a Solar Eclipse," June 1967.

229. Beyers, N. J., and B. T. Miers, "A Tidal Experiment in the Equatorial Stratosphere over Ascension Island (8S)", June 1967.

Miller, W. B., and H. Rachele, "On the Behavior of Derivative Processes," June 1967 230. Walters, Randall K., "Numerical Integration Methods for Ballistic Rocket Trajec-231. tory Simulation Programs," June 1967.

Hansen, Frank V., "A Diabatic Surface Boundary Layer Model," July 1967. 232.

Butler, Ralph L., and James K. Hall, "Comparison of Two Wind Measuring Sys-233. tems with the Contraves Photo-Theodolite," July 1967.

Webb, Willis L., "The Source of Atmospheric Electrification," June 1967.

- Hinds, B. D., "Radar Tracking Anomalies over an Arid Interior Basin," August 1967. 235.
- Christian, Larry O., "Radar Cross Sections for Totally Reflecting Spheres," August 236. 1967.
- D'Arcy, Edward M., "Theoretical Dispersion Analysis of the Aerobee 350," August 1967.
- Anon., "Technical Data Package for Rocket-Borne Temperature Sensor," August 238. 1967.
- Glass, Roy I., Roy L. Lamberth, and Ralph D. Reynolds, "A High Resolution Con-239. tinuous Pressure Sensor Modification for Radiosondes," August 1967.
- Low, Richard D. H., "Acoustic Measurement of Supersaturation in a Warm Cloud," 240. August 1967.
- 241. Rubio, Roberto, and Harold N. Ballard, "Time Response and Aerodynamic Heating of Atmospheric Temperature Sensing Elements," August 1967.
- Seagraves, Mary Ann B., "Theoretical Performance Characteristics and Wind Effects 242. for the Aerobee 150," August 1967.
- Duncan, Louis Dean, "Channel Capacity and Coding," August 1967. 243.
- Dunaway, G. L., and Mary Ann B. Seagraves, "Launcher Settings Versus Jack Settings for Aerobee 150 Launchers Launch Complex 35, White Sands 244. Missile Range, New Mexico," August 1967.
- Duncan, Louis D., and Bernard F. Engebos, "A Six-Degree-of-Freedom Digital Com-245. puter Program for Trajectory Simulation," October 1967.
- Rider, Laurence J., and Manuel Armendariz, "A Comparison of Simultaneous Wind 246. Profiles Derived from Smooth and Roughened Spheres," September 1967.
- Reynolds, Ralph D., Roy L. Lamberth, and Morton G. Wurtele, "Mountain Wave Theory vs Field Test Measurements," September 1967. 247.
- Lee, Robert P., "Probabilistic Model for Acoustic Sound Ranging," October 1967. 248.
- Williamson, L. Edwin, and Bruce Kennedy, "Meteorological Shell for Standard Artillery Pieces A Feasibility Study," October 1967. 249.
- Rohwer, Ralph H., "Acoustical, Meteorological and Seismic Data Report SOTRAN 250. V and VI," October 1967.
- Nordquist, Walter S., Jr., "A Study in Acoustic Direction Finding," November 1967. Nordquist, Walter S., Jr., "A Study of Acoustic Monitoring of the Gun Probe Sys-252.tem," November 1967.
- 253. Avara, E. P., and B. T. Miers, "A Data Reduction Technique for Meteorological Wind Data above 30 Kilometers," December 1967.
- Hansen, Frank V., "Predicting Diffusion of Atmospheric Contaminants by Considera-254. tion of Turbulent Characteristics of WSMR," January 1968.

UN	CLAS	SI	FIED	
_				

Security Classification							
DOCUMENT CONTROL DATA - R & D (Security classification of title, body of abstract and indexing annotation must be entered when the everall report is classified,							
1. ORIGINATING ACTIVITY (Corporate author)			CURITY CLASSIFICATION				
U. S. Army Electronics Command	1	Unclassified					
Fort Monmouth, New Jersey	!	M. SROUP					
·							
3. REPORT TITLE		<del></del>					
PREDICTING DIFFUSION OF ATMOSPHERIC CONTAMI	INANTS BY CON	SIDERATION	1 OF				
TURBULENT CHARACTERISTICS OF WSMR			i				
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)							
,							
5. AUTHOR(S) (First name, middle initial, last name)		<del></del>					
Frank V. Hansen			İ				
. REPORT DATE	74. TOTAL NO. OF	PAGES	78. NO. OF REFS				
January 1968	25		23				
M. CONTRACT OR GRANT NO.	M. ORIGINATOR'S REPORT NUMBER(S)						
b. PROJECT NO.	ECOM-5170						
	55. OTHER REPORT NOIS! (Any other numbers that may be essigned						
DA TASK 1T014501B53A-10	this report)						
	1		!				
10. DISTRIBUTION STATEMENT	<u> </u>						
			!				
Distribution of this report is unlimited							
11- SUPPLEMENTARY NOTES	12. SPONSORING M						
	U.S. Army Electronics Command						
	Atmospheric Sciences Laboratory White Sands Missile Range, New Mexico						
	White Sains	M155116 K	ange, New Mexico				
IS. ABSTRACT							
4							

The relationships between the turbulent intensities about the mean flow in the lowest layers of the atmosphere at White Sands Missile Range, New Mexico, and climatological data are discussed in terms of predicting the diffusion of contaminants into the atmosphere. If the turbulent fluctuations observed in the lower portion of the boundary layer are considered representative, then correlations with climatological summaries may be used as a rudimentary estimator for estimating the diffusion power of the atmosphere. A simple empirical diffusion model for general use is also given.

POR 4 A TY REPLACES DO PORM 1473, I JAN 64, WHICH IS

UNCLASSIFIED

Security Cassification

UNCLASSIFIED
Security Classification

LINK B KEY WORDS ROLE ROLE WT ROLE Diffusive Power Prediction Model Regression Curve
 Micrometeorological Study
 Turbulent Characteristics.

UNCLASSIFILD

Security Classification